

Coues, E. 1878. Birds of the Colorado Valley, U. S. G. of the Territories, Misc. Pub. No. 11. 807 pp.

DOMINICA

1836, 1837, 172.—*Bp. CGL.* 1838, 22.—*Atad. Syu.*  
—*Gosse, B. Jam.* 1847, 156.—*Denny, PZS.* 1847,  
p. CA. i. 1850, 307.—*Lawr. Ann. Lye. N. Y.* vi.  
Acad. vi. 1853, 398 (Ohio).—*Sallé, PZS.* 1857, 231  
i. 1860, 307 (Cuba).—*Albrecht, J. f. O.* 1862, 201  
, 91 (San Domingo).

474 (Cuba).—*Sci. PZS.* 1856, 291 (Mexico).—

i. 1855, 601.

73).  
1 (from Catesby, Brisson, &c.).—*Turt. SN.* i.

—*V. OAS.* ii. 1807, 45.—*Wils. AO.* ii. 1810, 64.  
—*Steph. Gen. Zool.* x. 1817, 679.—*V. Ency.*  
*Phila. Acad.* iv. 1824, 188.

1853, 310 (Wisconsin).

*ominicensis cinerea*, *Bris. Orn.* iii. 1763,  
L.).

E. 686, f. 1 (basis of *M. superciliosa* Bodd.

*ensis griseus*, *Bris. Orn.* iii. 1760, 563, n.

*guttare luteo*, *Cates. Car.* i. 1771, 62, pl. 62

n. 286.

783, 437, n. 31.

n. 37 (= *Coujaune*, *Buff. v.* 165; *PE.* 626, f.

, 292.

*urton*, ll. cc.

i. 110.

i. 1817, 191.

27.

45.—*V. Ency. Méth.* ii. 1823, 453.

ray Warbler, Yellow-throated Wood-

a

at. vii. 1873, 606.—*B. B. & R. NAB.* i. 1874,

names of this species. The first binomial,  
"de S. Domingue", after Sloane. The  
*sitis* Gm., are both from the "Coujaune"  
Latham. The fourth, *M. flavicollis* Gm.,

l, excepting *albilora* of Ridgway, which  
record of the bird from the Mississippi  
discriminate. Texas specimens I have  
irely white, and such seems to be the  
the range of the species.

herly. North regularly to Ohio,  
io Valley (*Wheaton et al.*); rarely  
ng latitudes. Washington, D. C.,  
*D. Scott*); Pennsylvania (*Turn-*  
*DeKay*). Connecticut (*cf. Coues*,

DENDRŒCA KIRTLANDI—D. PALMARUM 249

*Pr. Essex Inst.* v. 270; *Coues*, BNW. 66; and especially *Merriam*, *Tr. Conn.*  
*Acad.* iv. 17). West to Kansas (*Snow*) and Texas (*McCauley*). Breeds in its  
United States range at large. Winters in Florida, Mexico (Xalapa, Cordova,  
Colima), Central America, Guatemala, Yucatan, and various West India  
Islands (Cuba, San Domingo, Jamaica, where probably resident). Replaced  
in Porto Rico by *D. adelaida*, and in the Southern Middle Province of the  
United States by *D. gracia*.

It should be added, that Mr. Ridgway restricts the range of true *dominica*  
to the Atlantic States as far north as Washington in summer, and in winter  
to Cuba, San Domingo, and Jamaica, assigning to his var. *albilora* the follow-  
ing *habitat*:—"In summer, the Mississippi region of the United States, north  
to Lake Erie; common in South Illinois. In winter, and possibly all the  
year, in Mexico, south to Guatemala, Yucatan on the Atlantic, and Colima  
on the Pacific side."

*Dendroeca kirtlandi*.—Kirtland's Warbler.

*Sylvicola kirtlandii*, *Bd. Ann. Lye. N. Y.* v. 1852, 216, pl. 6 (Cleveland, Ohio, May, 1851, J.  
P. Kirtland).—*Zuchold, J. f. O.* 1854, 355 (copies descr.).—*Cass. Ill.* i. 1855, 278, pl. 47  
(from the original).

*Dendroica kirtlandii*, *Bd. BNA.* 1858, 286.—*Wheat. Ohio Agric. Rep.* for 1860, 1861, 374  
(ref. to the orig. spec. and to another from same locality, and to a possible third  
from Racine, Wis.).—*Bd. Rev. AB.* 1865, 206 (notifies a second specimen from sea  
near Bahamas, in Coll. S. Cabot, and report of third and fourth specimens in  
"Ohio Farmer" for June 9, 1860).

*Dendroeca kirtlandi*, *Sund. Oefv. K. Vet.-Akad. Förh.* iii. 1869, 617.—*Langdon, Cat. B.*  
*Cincinnati*, 1877, 6.

*Dendroeca kirtlandii*, *Coues. Key*, 1872, 104.

*Dendroica kirtlandi*, *B. B. & R. NAB.* i. 1874, 273, pl. 14, f. 5.

*Mniotilta kirtlandi*, *Gieb. Nomencl. Av.* ii. 1875, 603.

*Kirtland's Warbler*, *Authors*.

HAB.—Of the known specimens, four in number, three were taken in Ohio  
and one at sea between the Bahamas and Cuba. Supposed to have been seen  
in Wisconsin (Racine, *Hoy*).

*Dendroeca palmarum*.—Yellow Red-poll Warbler.

a. *palmarum*

*Motacilla palmarum*, *Gm. SN.* i. 1788, 951, n. 53 (San Domingo; based on the *Bimbelé ou*  
*fausse Linotte*, *Buff. v.* 330, and *Palm Warbler* of *Lath. Syn.* ii. pt. ii. 493, n. 131).—*Less.*  
*Tr. Orn.* 1831, 418.

*Sylvia palmarum*, *Lath. IO.* ii. 1790, 544, n. 136.—*V. OAS.* ii. 1807, 21, pl. 73.—*V. N. D. d'H. N.*  
xi. 1817, 163.—*Steph. Gen. Zool.* x. 1817, 607.—*V. Ency. Méth.* ii. 1823, 431, n. 42.—*Bp.*  
*AO.* ii. —, pl. 10, f. 2.—*Bp. Journ. Phila. Acad.* v. 1825, 23 (Florida).—*Bp. Ann.*  
*Lyc. N. Y.* ii. 1826, 78.—*D'Orb. Ois. Cuba*, 1839, 61, pl. 8.

*Sylvicola palmarum*, *Rich. Rep. Brit. Assoc. Adv. Sci.* for 1836, 1837, 172 (= *Bp. AO.* pl.  
10, f. 2).—*Sallé, PZS.* 1857, 231 (San Domingo).—*Bry. Pr. Bost. Soc.* vii. 1859, 110 (Baha-  
mas).—*Albrecht, J. f. O.* 1861, 53 (Bahamas).—*Bry. Pr. Bost. Soc.* xi. 1867, 91 (San  
Domingo).

*Mniotilta palmarum*, *Gray, G. of B.* i. 1848, 196 (= *Gm.*, and *Bp. AO.* pl. 10, f. 2).

*Dendroica palmarum*, *Bd. BNA.* 1858, 288.—*Barn. Smiths. Rep.* for 1860, 1861, 426.—*Gundl.*  
*J. f. O.* 1861, 326 (Cuba).—*Coues & Prent. Smiths. Rep.* for 1861, 1862, 408.—*Boardm. Pr.*  
*Bost. Soc.* ix. 1862, 123.—*Verr. Pr. Ess. Inst.* iii. 1862, 147.—*Allen, Pr. Ess. Inst.* iv. 1864,  
63.—*Bd. Rev. AB.* 1865, 207.—*Lawr. Ann. Lye. N. Y.* viii. 1866, 234.—*Trippe, Am. Nat.* ii.  
1868, 171.—*Turnb. B. E. Pa.* 1869, 25; *Phila. ed.* 18.—*Gundl. J. f. O.* 1872, 415 (Cuba).—  
*B. B. & R. NAB.* i. 1874, 273, pl. 14, f. 8.—*Brew. Pr. Bost. Soc.* xvii. 1875, 440.