





FINAL

Operational Range Assessment Program Phase I Qualitative Assessment Report Caswell Training Site, Maine

U.S. Army Operational Range Assessment Program Qualitative Operational Range Assessments

Prepared for:

U.S. Army Environmental Command and

U.S. Army Corps of Engineers Baltimore District





EXECUTIVE SUMMARY

The United States (U.S.) Army is conducting qualitative assessments at operational ranges to meet the requirements of Department of Defense policy and to support the U.S. Army Sustainable Range Program. The operational range qualitative assessment (hereinafter referred to as Phase I Assessment) is the first phase of the U.S. Army Operational Range Assessment Program (ORAP). This Phase I Assessment evaluates the operational range area at Caswell Training Site (TS) to assess whether further investigation is needed to determine if potential munitions constituents of concern (MCOC) are or could be migrating off-range at levels that may pose an unacceptable risk to human health or the environment. In conducting the Phase I Assessment, MCOC sources, potential off-range migration pathways, and potential off-range human and ecological receptors are evaluated as appropriate.

Caswell TS is a 1,049.08-acre facility that consists of three separate parcels (Caswell TS, Loring Prime Base Engineer Emergency Force [BEEF], and Loring Range) located in northeastern Maine, approximately four miles west of the U.S. / Canadian border. Initial use of the site began in 1947 when the U.S. Army's Strategic Air Command began the construction of Limestone Army Air Base. The Maine Army National Guard (MEARNG) began using the site in the mid-1960s and was granted federal license in the 1970s. Loring Air Force Base was closed under Base Realignment and Closure in the early 1990s; the MEARNG requested the Loring Prime BEEF Area and Loring Range at that time and was granted license in 1999. The Army Range Inventory Database-Geodatabase (2007) identifies five operational range areas at Caswell TS consisting of maneuver and training areas and firing ranges encompassing the entire 1,049.08 acres.

Primarily, MCOC sources identified at Caswell TS consist of current and historical small caliber firing points and impact berms. In general, MCOC from source areas potentially impact soil (e.g., impact berms) and surface water / sediment (e.g., deposition into streams and wetlands).

MCOC can be released to groundwater (down gradient), surface water / sediment (downstream), or the food chain via a variety of release mechanisms. Release mechanisms for soil may include erosion and runoff to nearby streams and wetlands or leaching/infiltration to surficial deposits or the bedrock aquifer. Once potential MCOC are deposited in surface water / sediment, they have the potential to migrate downstream and/or be taken up by aquatic plants or animals. Release mechanisms for surface water / sediment are natural stream flow and sediment transport. Once potential MCOC leach/infiltrate into the surficial deposits or bedrock aquifer they have the potential to discharge into streams and wetlands or be taken up through groundwater wells. Surface water and groundwater at Caswell TS generally flow to the south and southeast, with the exception of Loring Prime BEEF where surface and groundwater apparently flow to the southwest.

The five operational ranges at Caswell TS are categorized as Unlikely.

Unlikely – Five-Year Review

The five ranges at Caswell TS are categorized as Unlikely, totaling 1,049.08 acres. These ranges consist of maneuver and training areas and firing ranges. Ranges where, based upon a review of readily available information, there is sufficient evidence to show that there are no known releases or source-receptor interactions off-range that could present an unacceptable risk to human health or the environment are categorized as Unlikely. Ranges categorized as Unlikely are required to be re-

evaluated at least every five years. Re-evaluation may occur sooner if significant changes (e.g., change in range operations or site conditions, regulatory changes) occur that affect determinations made during this Phase I Assessment.

Table ES-1 summarizes the Phase I Assessment findings.

Table ES-1: Summary of Findings and Conclusions for Caswell Training Site

Catalana	Total Number of Ranges and	S(-)	D. (b(a)	Human	Ender'n Derman	Cardada and D. Carda
Category Unlikely	5 operational ranges; 1,049.08 acres	No source— limited or no military munitions use	Pathway(s) Not evalua	Receptors ated (no source wa	Ecological Receptors as identified)	Re-evaluate during the five-year review. No source was identified.
		Historical small arms firing points and impact berm	Wetlands and intermittent streams	Off-range residents downstream and down gradient	Sensitive environments (i.e., Aroostook National Wildlife Refuge, upland sandpiper, and wetlands)	Re-evaluate during the five-year review. Based on on-range wetland characteristics and the level of vegetation at the site, limited potential for off-range MCOC migration was identified.
		Active small arms firing points and impact berm	Surficial deposits and bedrock aquifer			Re-evaluate during the five-year review. Based on lead removal activities and the characteristics of the surficial deposits and bedrock aquifer, limited potential for off-range MCOC migration was identified.

ABBREVIATIONS/ACRONYMS

ANWR	Aroostook National Wildlife Refuge	
ARID-GEO	Army Range Inventory Database-Geodatabase	
BEEF	Base Engineer Emergency Force	
BRAC	Base Realignment and Closure	
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act	
CERCLA	Conceptual Site Model	
	<u> </u>	
DNT	Dinitrotoluene	
DoD	Department of Defense	
DODI	Department of Defense Instruction	
E	Ecological receptors identified. (This refers to range grouping; pathway designation always precedes E designation.)	
ERM	Environmental Resources Management	
GW	Groundwater pathway identified. (This refers to range grouping; M	
011	designation always precedes GW designation.)	
Н	Human receptors identified. (This refers to range grouping; pathway	
11	designation always precedes H designation.)	
ITAM	Integrated Training Area Management	
LAFB	Loring Air Force Base	
LS	Limited Source	
M	Munitions used. (This refers to range grouping; M designation always	
141	precedes applicable pathway.)	
MCOC	Munitions Constituents of Concern	
MEARNG	Maine Army National Guard	
MEDEP	Maine Department of Environmental Protection	
MEDWP	Maine Drinking Water Program	
MEGIS	Maine Office of Geographic Information Systems	
NG	Nitroglycerin	
NOAA	National Oceanic and Atmospheric Administration	
NRCS	Natural Resources Conservation Service	
ORAP	Operational Range Assessment Program	
RFMSS	Range Facility Management Support System	
SW	Surface water pathway identified. (This refers to range grouping; M	
5 **	designation always precedes SW designation.)	
TS	Training Site	
U.S.	United States	
USACE	United States United States Army Corps of Engineers	
USACHPPM	United States Army Corps of Engineers United States Army Center for Health Promotion and Preventive Medicine	
USAEC	United States Army Environmental Command	
USAF	United States Air Force	
USDA	United States Department of Agriculture	
USEPA	United States Environmental Protection Agency	
USFWS	United States Fish and Wildlife Service	
USGS	United States Geological Survey	
°F	Degrees Fahrenheit	
Г	Degrees Pattietitiett	






Operational Range Assessment Program Phase I Qualitative Assessment Caswell Training Site, ME



Figure 1-1 General Caswell Training Site Location



Installation

Installation Boundary

Data Sources: ARID-GEO 2007, ESRI Canada Map 2006, ESRI StreetMap USA 2005

Date:.....July 2008
Prepared By: EA Engineering, Science & Technology
Prepared For:....U.S. Army
Contract Number:....W912DR-05-D-0008