```
FIELD VERIFICATION REPORT
                                                         Submitted to Rare Birds Comm
                                                         Kingston Field Naturalists
                                                         Kingston, Ontario
          Date of Sighting: 1981 Sept. 9 about 0900, seen clearly for about one min
1.
2.
          Location:
                      Morton (about 25 mi. NE of Kingston, Ontario)
3.
          Habitat:
                      White Pine stand near road with edge of deciduous bushes and
                      saplings.
          Weather:
                      Broken cloud, sun behind observer.
5.
          Optical aids: 8 x 40 Binocs in excellent condition and bird viewer at
                      very close range at about 10 paces, allowed close approach.
6.
          Field Marks:
              a. General
                      Chickadee-sized bird feeding in pines, dark above with wing
                 bars and bright yellow beneath. The unhooked small dark bill marked
                 bird as a warbler. It wagged tail continuously much like Phoebe.
              b. Detail
                     (i) Size of black-capped Chickadee feeding in same trees but
                 warbler.chunkier.
                    (ii) Underparts - bright unstreaked yellow throat, breast and
                 belly becoming white near vent and under tail coverts. Flanks,
                 sides up to shoulder showed light black streaking.
                   (iii) Head and nape - gray.
                    (iv) Face - dark with distinctive white eye ring that appeared
                 interrupted or broken by black extending from bill to eye. No
                 yellow was present on face and black in front of eye to bill fairly
                 extensive, some black in ear region.
                     (v) Upper-parts - white wing bars (2) contrasting with dark
                 wings. Do not recall detail of the back since book consulted and
                 upon relooking at bird, it flew to an inaccessible area of wood.
                    (vi) Tail wagging - continuous; up and down motion.
              c. Behaviour
                    - gathering insects/food from pine needles and branches, but
                 nothing distinctive noted about its method of feeding. No calls or
                 song was heard.
          Species - Kirtland's Warbler (1) male (black on face).
7.
```

```
i) Prairie - ruled out because it shows yellow in face in adult
                    plumage. The young Prairie shows a white throat.
                ii) Palm - ruled out because it also shows yellow in face, red/wine/
                    brown cap and some breast streaking. Its vent and under tail
                    coverts are yellow.
                     Marg Hendrick and Tim Hendrick.
9.
          Observers:
          Previous experience with species: One sighting with Dr. Paul Aird
10.
                    within last few years on Kirtland's survey in Quebec.
                                           II . YS O G W W B ST O L
          Comments: (a) Missed seeing detail on the back as bird fed overhead, which
11.
                    is only field mark missing. Back was dark however.
                    (b) Above information given to Ron Weir by telephone on Sept. 9
                    for Rare Birds Committee.
                                               West floor change are
                                         - Estadorial (11)
                                         en at his putaosid villa.
                                        taes uneto stoutden
```

There are 3 tail-wagging warblers of North America, viz. Kirtland's,

com bus fresh (iti)

com bus fresh (iti)

com on best (iti)

com on best (iti)

com on best (iti)

27780-7901U (v1981 Sept.10

Mag of the second

and the first state of

The part of the part of the control of the control

pathons from (10)

w dathertro for

Similar confusing species eliminated by description.

8.

Prairie and Palm.